

Specific heat and validity of quasiparticle approximation in the half-filled Landau level

Yong Baek Kim

Bell Laboratories, 600 Mountain Avenue, Murray Hill, NJ 07974

Patrick A. Lee

Department of Physics, Massachusetts Institute of Technology, Cambridge, MA 02139

Abstract

We calculate the specific heat of composite fermion system in the half-filled Landau level. Two different methods are used to examine validity of the quasiparticle approximation when the two-body interaction is given by $V(q) = V_0/q^{2-\eta}$ ($1 \leq \eta \leq 2$). The singular part of the specific heat is calculated from the free energy of the gauge field, which is compared with the specific heat calculated from the quasiparticle approximation via the singular self-energy correction due to the gauge field fluctuations. It turns out that two results are in general different and they coincide only for the case of the Coulomb interaction ($\eta = 1$). This result supports the fact that the quasiparticle approximation is valid only for the case of the Coulomb interaction. It is emphasized that this result is obtained by looking at a gauge-invariant quantity – the specific heat.

Pacs numbers: 73.40.Hm, 71.10.Pm, 11.15.-q

The appearance of the unexpected metallic state at the filling fraction $\nu = 1/2$ in the fractional quantum Hall regime has stimulated a lot of activities in theory^{1-9,10-16} and experiment¹⁷⁻²⁵. The similarity between the phenomena around $\nu = 1/2$ in high magnetic fields and those of electrons in weak magnetic fields is successfully explained by the concept of composite fermions¹. A composite fermion is obtained by attaching even number of flux quanta to an electron¹⁻³. In particular, in the half-filled Landau level, one attaches two flux quanta to an electron to make a composite fermion. At the mean field level, if one takes the average of the statistical magnetic field coming from the attached magnetic flux, the composite fermions at $\nu = 1/2$ feel zero magnetic field due to the cancellation between the external magnetic field and the averaged statistical magnetic field³. As a result, the mean field ground state is a filled Fermi-sea with a well-defined Fermi wave vector k_F ³. There exist several experiments which demonstrate the existence of a well-defined Fermi wave vector at $\nu = 1/2$ ²²⁻²⁵.

Note that the fluctuations of the statistical magnetic field correspond to the density fluctuations due to the fact that the statistical flux quanta are attached to each electron. Therefore, in the mean field approximation, the strong density fluctuations are ignored. This implies that the fluctuations about the mean field state can be very important and are basically gauge field fluctuations. The above arguments also suggest that the effects of the two-particle interaction $v(\mathbf{q}) = V_0/q^{2-\eta}$ ($1 \leq \eta \leq 2$) on the gauge field fluctuations should be examined carefully. In fact, the gauge field fluctuations become more singular as the interaction range becomes shorter (larger η) because the longer range interaction (smaller η) suppresses the density fluctuations more effectively^{3,5}. Thus, it is important to examine the stability of the mean field Fermi-liquid state against the gauge field fluctuations.

In order to study systematically the effects of the gauge field fluctuations, Halperin, Lee, and Read (HLR) used the fermionic Chern-Simons gauge theory to develop a theory of the metallic state at $\nu = 1/2$ ³. In the random phase approximation which becomes exact in the large N limit (N is the number of species of fermions), the most singular correction to the self-energy comes from the transverse part of the gauge field, which is given by³

$$D_{11}(\mathbf{q}, \nu) = \frac{1}{-i\gamma\nu/q + \chi q^\eta} , \quad (1)$$

where $\gamma = 2n_e/k_F$, $\chi = \frac{1}{24\pi m} + \frac{V_0}{(2\pi\phi)^2}$ for $\eta = 2$, and $\chi = \frac{V_0}{(2\pi\phi)^2}$ for $1 \leq \eta < 2$. The correction to the retarded self-energy in the lowest order perturbation theory is given by^{3,5,26}, for $1 < \eta \leq 2$,

$$\Sigma^R(\mathbf{k}, \omega) = [\text{Re}\lambda + i \text{Im}\lambda \text{sgn}(\omega)] |\omega|^{\frac{2}{1+\eta}} , \quad (2)$$

where

$$\lambda = \frac{v_F e^{i\frac{\pi}{2}(\frac{\eta-1}{\eta+1})}}{4\pi \sin\left(\frac{2\pi}{1+\eta}\right) \gamma^{\frac{\eta-1}{\eta+1}} \chi^{\frac{2}{1+\eta}}} . \quad (3)$$

For $\eta = 1$, we have

$$\Sigma^R(\mathbf{k}, \omega) = \frac{v_F}{4\pi^2\chi} |\omega| \left[\ln\left(\frac{4k_F^2\chi}{\gamma|\omega|}\right) + i\frac{\pi}{2}\text{sgn}(\omega) \right] . \quad (4)$$

Therefore, the usual Landau criterion for the quasiparticle is violated in the case of $1 < \eta \leq 2$ and the case of $\eta = 1$ shows the marginal Fermi liquid behavior.

In Ref.³, an attempt was made to construct a renormalized quasiparticle theory using the above singular self-energy correction even though there is no well-defined quasiparticle in the usual sense of Landau-Fermi-liquid theory. It was assumed that there exists a well defined Fermi wave vector $k_F = \sqrt{4\pi n_e}$ and, for $|\mathbf{k}| \approx k_F$, there exist quasiparticle excitations with the energy spectrum³, for $1 < \eta \leq 2$,

$$E(\mathbf{k}) = (|\xi_{\mathbf{k}}|/\text{Re}\lambda)^{\frac{1+\eta}{2}} \text{sgn}(\xi_{\mathbf{k}}) , \quad (5)$$

where $\xi_{\mathbf{k}} = \frac{k^2}{2m} - \mu$. For $\eta = 1$, the quasiparticle spectrum becomes

$$E(\mathbf{k}) = \frac{4\pi^2\chi}{v_F} \xi_{\mathbf{k}} \ln\left(\frac{k_F^2 v_F}{\pi^2 \gamma |\xi_{\mathbf{k}}|}\right) . \quad (6)$$

Note that these quasiparticle spectra are obtained from the real part of the retarded self-energy.

Recently we derived and studied the quantum Boltzmann equation (QBE) of the composite fermions interacting with the gauge field⁹. It was emphasized that a generalized

QBE can be constructed even though there is no well defined Landau-quasiparticle⁹. In particular, the collision integral of the QBE becomes relatively unimportant only when the two-particle interaction is given by the Coulomb interaction ($\eta = 1$)⁹. In this case, the QBE is equivalent to the usual QBE of Landau-Fermi-liquid theory. For the case of Coulomb interaction, Stern and Halperin⁸ showed that the usual Landau-Fermi-liquid theory can be successfully applied, which is consistent with the construction of the QBE mentioned above. Also Kwon, Houghton, and Maston¹³ argued that the higher dimensional bosonization can be consistently performed only for the case of the Coulomb interaction ($\eta = 1$). This result may imply that the case of the Coulomb interaction can be described by the usual Landau-Fermi-liquid theory.

In this paper, we want to examine the range of the applicability of the quasiparticle approximation in relation to the Landau-Fermi-liquid theory. In particular, we address this issue by looking at a gauge-invariant physical quantity – specific heat. According to the standard procedure, the total specific heat of the system can be calculated as the sum of the contributions from the free fermion and the gauge field³. The free fermion part gives linear temperature dependence while the gauge field part gives more singular non-linear temperature dependence in the low temperature limit. In the quasiparticle approximation, the singular correction to the specific heat could be considered as coming from the singular mass renormalization of the fermions. In fact, both of the methods give the same temperature dependence for the singular part of the specific heat. The question is whether they are exactly the same in the low temperature limit. Thus, as a direct test of the quasiparticle approximation, we calculate the most singular part of the specific heat using two different methods. One is to use the dispersion relation of the renormalized quasiparticle in the quasiparticle approximation. The other way is to calculate the specific heat from the free energy of the gauge field. The latter method is supposed to give the correct answer because the latter one is gauge-invariant while the former is not. It turns out that the two results are in general different and they agree with each other only for the case of the Coulomb interaction. This result supports the fact that the quasiparticle approximation can be safely

used only for the case of the Coulomb interaction. We want to emphasize that this result is obtained by looking at a gauge-invariant quantity – the specific heat.

First, let us evaluate the specific heat using the quasiparticle approximation. Using the dispersion relation given by Eqs. 5 and 6, the expectation value of the energy can be obtained as

$$\langle E \rangle = \sum_{\mathbf{k}} E(\mathbf{k}) f(\mathbf{k}) , \quad (7)$$

where $f(\mathbf{k}) = 1/(e^{E(\mathbf{k})/T} + 1)$. The specific heat can be evaluated from $C_{\text{qp}} = \frac{\partial \langle E \rangle}{\partial T}$:

$$\frac{\partial \langle E \rangle}{\partial T} = \frac{m}{4\pi} \int_0^\infty d\xi_{\mathbf{k}} \frac{E^2(\mathbf{k})/T^2}{[\cosh(E(\mathbf{k})/2T)]^2} . \quad (8)$$

For $1 < \eta \leq 2$, it is given by

$$C_{\text{qp}}(T) = \alpha_{\text{qp}}(\eta) T^{\frac{2}{1+\eta}} , \quad (9)$$

where

$$\alpha_{\text{qp}}(\eta) = \frac{1 - 2^{-\frac{2}{1+\eta}}}{2\pi(1+\eta)} \frac{1}{\sin\left(\frac{\pi}{2} \frac{\eta-1}{\eta+1}\right)} \Gamma\left(\frac{2\eta+4}{\eta+1}\right) \zeta\left(\frac{\eta+3}{\eta+1}\right) \left(\frac{\gamma}{\chi}\right)^{\frac{2}{1+\eta}} . \quad (10)$$

In the case of $\eta = 1$, we have

$$C_{\text{qp}}(T) = \frac{1}{12} \frac{\gamma}{\chi} T \ln\left(\frac{4k_F^2 \chi}{\gamma T}\right) . \quad (11)$$

Now we calculate the specific heat from the free energy of the gauge field:

$$F_{\text{g}} = \int_{-\infty}^{\infty} \frac{d\nu}{2\pi} \int \frac{d^2 q}{(2\pi)^2} \frac{1}{e^{\nu/T} - 1} \arctan\left(\frac{\text{Im} D_{11}^{-1}}{\text{Re} D_{11}^{-1}}\right) . \quad (12)$$

The contribution of the gauge field to the entropy can be obtained as

$$S_{\text{g}} = -\frac{\partial F_{\text{g}}}{\partial T} = \int_{-\infty}^{\infty} \frac{d\nu}{2\pi} \int_0^\infty \frac{qdq}{2\pi} \frac{\nu/T^2}{(e^{\nu/2T} - e^{-\nu/2T})^2} \arctan\left(\frac{\gamma}{\chi} \frac{\nu}{q^{1+\eta}}\right) . \quad (13)$$

Thus, for $1 < \eta \leq 2$, the specific heat $C_{\text{g}} = T \frac{\partial S_{\text{g}}}{\partial T}$ is given by

$$C_{\text{g}}(T) = \alpha_{\text{g}}(\eta) T^{\frac{2}{1+\eta}} , \quad (14)$$

where

$$\alpha_g(\eta) = \frac{1}{4\pi(1+\eta)} \frac{1}{\sin\left(\frac{\pi}{2}\frac{\eta-1}{\eta+1}\right)} \Gamma\left(\frac{2\eta+4}{\eta+1}\right) \zeta\left(\frac{\eta+3}{\eta+1}\right) \left(\frac{\gamma}{\chi}\right)^{\frac{2}{1+\eta}}. \quad (15)$$

In the case of $\eta = 1$, the specific heat becomes the same as Eq. 11.

Note that $C_g(T)$ is the correct answer for the singular part of the specific heat because it is calculated in a gauge-invariant way. Note also that $C_g(T)$ agrees with the result $C_{qp}(T)$ of the quasiparticle approximation only for the case of $\eta = 1$. The underlying reason for this behavior is as follows. In the quasiparticle approximation, only the real part of the self-energy is used to get the spectrum of the quasiparticle. However, both of the real and imaginary parts of the self-energy contribute to the specific heat, *i.e.*, one has to also take into account the information about the imaginary part of the self-energy. The sum of these contributions are basically the same as the specific heat calculated from the free energy of the gauge field⁷. For the case of the Coulomb interaction, the imaginary part of the self-energy is logarithmically smaller than the real part of the self-energy so that the real part of the self-energy is sufficient to get the correct answer for the most singular part of the specific heat. This result supports the previous conclusion that the quasiparticle approximation can be justified only when the two-particle interaction is given by the Coulomb interaction ($\eta = 1$).

We would like to thank B. I. Halperin, F. D. M. Haldane and X.-G. Wen for helpful discussions about this and related issues in relation to the half-filled Landau level. PAL is supported by NSF grant No. DMR-9523361.

REFERENCES

- ¹ J. K. Jain, Phys. Rev. Lett. **63**, 199 (1989); Phys. Rev. B **41**, 7653 (1990); Adv. Phys. **41**, 105 (1992).
- ² A. Lopez and E. Fradkin, Phys. Rev. B **44**, 5246 (1991); Phys. Rev. Lett. **69**, 2126 (1992).
- ³ B. I. Halperin, P. A. Lee, and N. Read, Phys. Rev. B **47**, 7312 (1993).
- ⁴ V. Kalmeyer and S. C. Zhang, Phys. Rev. B **46**, 9889 (1992).
- ⁵ Y. B. Kim, A. Furusaki, X.-G. Wen, and P. A. Lee, Phys. Rev. B **50**, 17917 (1994).
- ⁶ B. L. Altshuler, L. B. Ioffe, and A. J. Millis, Phys. Rev. B **50**, 14048 (1994).
- ⁷ Y. B. Kim, P. A. Lee, X.-G. Wen, and P. C. E. Stamp, Phys. Rev. B **51**, 10779 (1995).
- ⁸ A. Stern and B. I. Halperin, Phys. Rev. B **52**, 5890 (1995).
- ⁹ Y. B. Kim, P. A. Lee, and X.-G. Wen, Phys. Rev. B **52**, 17275 (1995).
- ¹⁰ D. V. Khveshchenko and P. C. E. Stamp, Phys. Rev. Lett. **71**, 2118 (1993); Phys. Rev. B **49**, 5227 (1994).
- ¹¹ J. Polchinski, Nucl. Phys. **B 422**, 617 (1994).
- ¹² Junwu Gan and Eugene Wong, Phys. Rev. Lett. **71**, 4226 (1994).
- ¹³ H. -J. Kwon, A. Houghton, and J. B. Marston, Phys. Rev. Lett. **73**, 284 (1994); Phys. Rev. B **52**, 8002 (1995).
- ¹⁴ C. Nayak, and F. Wilczek, Nucl. Phys. **B 417**, 359 (1994); Nucl. Phys. **B 430**, 534 (1994).
- ¹⁵ S. He, P. M. Platzman, and B. I. Halperin, Phys. Rev. Lett. **71**, 777 (1993); Y. Hatsugai, P.-A. Bares, and X.-G. Wen, Phys. Rev. Lett. **71**, 424 (1993); Y. B. Kim and X.-G. Wen, Phys. Rev. B **50**, 8078 (1994).
- ¹⁶ S. H. Simon and B. I. Halperin, Phys. Rev. B **48**, 17368 (1993); Phys. Rev. B **50**, 1807

- (1994); Song He, S. H. Simon, and B. I. Halperin, Phys. Rev. B **50**, 1823 (1994).
- ¹⁷ H. W. Jiang *et al.*, Phys. Rev. B **40**, 12013 (1989).
- ¹⁸ R. R. Du *et al.*, Phys. Rev. Lett. **70**, 2944 (1993).
- ¹⁹ D. R. Leadley *et al.*, Phys. Rev. Lett. **72**, 1906 (1994).
- ²⁰ R. R. Du *et al.*, Phys. Rev. Lett. **73**, 3274 (1994).
- ²¹ H. C. Manoharan, M. Shayegan, and S. J. Klepper *et al.*, Phys. Rev. Lett. **73**, 3270 (1994).
- ²² R. L. Willet *et al.*, Phys. Rev. Lett. **71**, 3846 (1993); R. L. Willet *et al.*, Phys. Rev. B **47**, 7344 (1993).
- ²³ W. Kang *et al.*, Phys. Rev. Lett. **71**, 3850 (1993).
- ²⁴ V. J. Goldman, B. Su, and J. K. Jain, Phys. Rev. Lett. **72**, 2065 (1994).
- ²⁵ R. L. Willet, K. W. West, and L. N. Pfeiffer, Phys. Rev. Lett. **75**, 2988 (1995).
- ²⁶ N. Nagaosa and P. A. Lee, Phys. Rev. Lett. **64**, 2550 (1990); P. A. Lee and N. Nagaosa, Phys. Rev. B **46**, 5621 (1992).